# CCDC61 Rabbit pAb

**Catalog NO.: A10341**

## Basic Information

### Observed MW
57kDa

### Calculated MW
57kDa

### Category
Polyclonal Antibodies

### Applications
WB, ELISA

### Cross-Reactivity
Human

### Conjugation
Unconjugated

## Background

Enables identical protein binding activity and microtubule binding activity. Involved in centriole assembly and mitotic spindle assembly. Located in centriolar satellite and ciliary basal body. Part of centriolar subdistal appendage.

## Recommended Dilutions

**WB** 1:500 - 1:2000

### ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID
729440

### Swiss Prot
Q9Y6R9

### Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms
VFL3; hVFL3; CCDC61

## Contact

## Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


**Source**
Rabbit

**Isotype**
IgG

**Purification**
Affinity purification

**Storage**
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data




Western blot analysis of lysates from Jurkat cells, using CCDC61 Rabbit pAb (A10341) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.