Cardiac troponin T (TNNT2) Rabbit pAb

Catalog NO.: A1126SP **4 Publications**

Basic Information

Observed MW
40 kDa

Calculated MW
31-36 kDa

Category
Polyclonal Antibodies

Applications
WB, IF-P, IHC-P, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

This gene encodes the cardiac isoform of troponin T. The encoded protein is the tropomyosin-binding subunit of the troponin complex, which is located on the thin filament of striated muscles and regulates muscle contraction in response to alterations in intracellular calcium ion concentration. Mutations in this gene have been associated with familial hypertrophic cardiomyopathy as well as with dilated cardiomyopathy.

Recommended Dilutions

WB	1:2000 - 1:6000
IF-P	1:200 - 1:2000
IHC-P	1:5000 - 1:20000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (⩾1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
7139

Swiss Prot
P45379

Immunogen
This information is considered to be commercially sensitive.

Synonyms
CMH2; RCM3; TnTC; cTnT; CMD1D; CMPD2; LVNC6; Cardiac troponin T (TNNT2)

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.


Validation Data





Western blot analysis of various lysates using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP) at 1:1000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): Mouse skeletal muscle.
Exposure time: 10 s.



Western blot analysis of various lysates using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP) at 1:6000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Negative control (NC): 293T.
Exposure time: 60 s.








Immunohistochemistry analysis of paraffin-embedded Mouse heart tissue using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP) at a dilution of 1:6000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Rat heart tissue using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP) at a dilution of 1:6000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.


Confocal imaging of paraffin-embedded Human heart tissue using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit **https://abclonal.com** for a complete list of recommended products.

Validation Data



Confocal imaging of paraffin-embedded Mouse heart tissue using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.



Confocal imaging of paraffin-embedded Rat heart tissue using Cardiac troponin T (TNNT2) Rabbit pAb (A1126SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.