JADE1 Rabbit pAb

Catalog NO.: A12579

Basic Information

Observed MW

Calculated MW
96kDa

Category
Polyclonal Antibodies

Applications
IF/ICC, ELISA

Cross-Reactivity
Human, Mouse

Conjugation
Unconjugated

Background

Enables transcription coactivator activity. Involved in histone acetylation and negative regulation of canonical Wnt signaling pathway. Acts upstream of or within negative regulation of G1/S transition of mitotic cell cycle. Located in several cellular components, including ciliary basal body; cytosol; and nuclear speck. Part of histone acetyltransferase complex.

Recommended Dilutions

IF/ICC	1:50 - 1:100

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
79960

Swiss Prot
Q6IE81

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
PHF17; JADE1

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunofluorescence analysis of U2OS cells using JADE1 Rabbit pAb (A12579). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.