

LIFR Rabbit pAb

Catalog NO.: A12783

Basic Information

Observed MW

190kDa

Calculated MW

124kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse

Conjugation

Unconjugated

Background

This gene encodes a protein that belongs to the type I cytokine receptor family. This protein combines with a high-affinity converter subunit, gp130, to form a receptor complex that mediates the action of the leukemia inhibitory factor, a polyfunctional cytokine that is involved in cellular differentiation, proliferation and survival in the adult and the embryo. Mutations in this gene cause Schwartz-Jampel syndrome type 2, a disease belonging to the group of the bent-bone dysplasias. A translocation that involves the promoter of this gene, t(5;8)(p13;q12) with the pleiomorphic adenoma gene 1, is associated with salivary gland pleiomorphic adenoma, a common type of benign epithelial tumor of the salivary gland. Multiple splice variants encoding two different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

3977

Swiss Prot

P42702

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

SWS; SJS2; STWS; CD118; LIF-R; LIFR


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of lysates from mouse skeletal muscle, using LIFR Rabbit pAb (A12783) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 30s.

