

MYH6/ α -MHC Rabbit pAb

Catalog NO.: A12964SP **13 Publications**

Basic Information

Observed MW

250 kDa

Calculated MW

224 kDa

Category

Polyclonal Antibodies

Applications

WB, IF-P, IHC-P, ELISA

Cross-Reactivity

Mouse, Rat

Conjugation

Unconjugated

Background

Cardiac muscle myosin is a hexamer consisting of two heavy chain subunits, two light chain subunits, and two regulatory subunits. This gene encodes the alpha heavy chain subunit of cardiac myosin. The gene is located approximately 4kb downstream of the gene encoding the beta heavy chain subunit of cardiac myosin. Mutations in this gene cause familial hypertrophic cardiomyopathy and atrial septal defect 3.

Recommended Dilutions

WB 1:2000 - 1:20000

IF-P 1:100 - 1:200

IHC-P 1:1000 - 1:4000

ELISA

Recommended starting concentration is 1 μ g/mL.
Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions ($\geq$ 1:10000)

Immunogen Information

Gene ID	Swiss Prot
4624	P13533

Immunogen

This information is considered to be commercially sensitive.

Synonyms

ASD3; MYHC; SSS3; CMH14; MYHCA; CMD1EE; alpha-MHC; MYH6/ α -MHC

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.

Validation Data



Western blot analysis of various lysates using MYH6/α-MHC Rabbit pAb (A12964SP) at 1:20000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Negative control (NC): Mouse kidney.

Exposure time: 20 s.

