

U2AF1 Rabbit pAb

Catalog NO.: A13166 **2 Publications**

Basic Information

Observed MW

38kDa

Calculated MW

28kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

This gene belongs to the splicing factor SR family of genes. U2 auxiliary factor, comprising a large and a small subunit, is a non-snRNP protein required for the binding of U2 snRNP to the pre-mRNA branch site. This gene encodes the small subunit which plays a critical role in both constitutive and enhancer-dependent RNA splicing by directly mediating interactions between the large subunit and proteins bound to the enhancers. Alternatively spliced transcript variants encoding different isoforms have been identified.

Recommended Dilutions

WB 1:1000 - 1:3000

IHC-P 1:100 - 1:200

IP
0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA
Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID	Swiss Prot
7307	Q01081

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.


Synonyms
RN; FP793; U2AF35; U2AFBP; RNU2AF1; U2AF1

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with thimerosal or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.		

Validation Data



Western blot analysis of various lysates using U2AF1 Rabbit pAb (A13166) at 1:3000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg U2AF1 antibody (A13166). Western blot was performed from the immunoprecipitate using U2AF1 antibody (A13166) at a dilution of 1:1000.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using U2AF1 Rabbit pAb (A13166) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.