

UPF2 Rabbit pAb

Catalog NO.: A13411 **1 Publications**

Basic Information

Observed MW

170kDa

Calculated MW

148kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, IP, ELISA

Cross-Reactivity

Human, Rat

Conjugation

Unconjugated

Background

This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located in the perinuclear area. It interacts with translation release factors and the proteins that are functional homologs of yeast Upf1p and Upf3p. Two splice variants have been found for this gene; both variants encode the same protein.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID

26019

Swiss Prot

Q9HAU5

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

HUPF2; RENT2; smg-3; UPF2


Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Western blot analysis of various lysates, using UPF2 Rabbit pAb (A13411) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.



Immunoprecipitation analysis of 300ug extracts of 293T cells using 3ug UPF2 Rabbit pAb (A13411 1:200). Western blot was performed from the immunoprecipitate using UPF2 Rabbit pAb (A7091) at a dilution of 1:1000.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using UPF2 Rabbit pAb (A13411) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.