

FA2H Rabbit pAb

Catalog NO.: A13874

Basic Information

Observed MW

40kDa

Calculated MW

43kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse, Rat

Conjugation

Unconjugated

Background

This gene encodes a protein that catalyzes the synthesis of 2-hydroxysphingolipids, a subset of sphingolipids that contain 2-hydroxy fatty acids. Sphingolipids play roles in many cellular processes and their structural diversity arises from modification of the hydrophobic ceramide moiety, such as by 2-hydroxylation of the N-acyl chain, and the existence of many different head groups. Mutations in this gene have been associated with leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

79152

Swiss Prot

Q7L5A8

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

FAAH; FAH1; SCS7; SPG35; FAXDC1; FA2H

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of various lysates using FA2H Rabbit pAb (A13874) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.

