

# NFE2L1 Rabbit pAb

Catalog NO.: A14753 **2 Publications**

## Basic Information

### Observed MW

120kDa

### Calculated MW

85kDa

### Category

Polyclonal Antibodies

### Applications

WB, IHC-P, IF/ICC, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

This gene encodes a protein that is involved in globin gene expression in erythrocytes. Confusion has occurred in bibliographic databases due to the shared symbol of NRF1 for this gene, NFE2L1, and for "nuclear respiratory factor 1" which has an official symbol of NRF1.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

| Gene ID | Swiss Prot |
|---------|------------|
| 4779    | Q14494     |

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

NRF1; NRF-1; TCF11; LCR-F1; NFE2L1


## Contact

## Product Information


|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                         |                |                       |
|---------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                           | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                  | IgG            | Affinity purification |
| <b>Storage</b>                                          |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.             |                |                       |
| Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3. |                |                       |


Validation Data




Western blot analysis of various lysates using NFE2L1 Rabbit pAb (A14753) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.




Immunohistochemistry analysis of paraffin-embedded Rat heart using NFE2L1 Rabbit pAb (A14753) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta using NFE2L1 Rabbit pAb (A14753) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using NFE2L1 Rabbit pAb (A14753) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of HeLa cells using NFE2L1  
Rabbit pAb (A14753) at dilution of 1:100. Secondary  
antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)  
(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.