

UBAC2 Rabbit pAb

Catalog NO.: A15977

Basic Information

Observed MW

39kDa

Calculated MW

39kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse

Conjugation

Unconjugated

Background

Involved in negative regulation of canonical Wnt signaling pathway and negative regulation of retrograde protein transport, ER to cytosol. Acts upstream of or within protein localization to endoplasmic reticulum. Located in endoplasmic reticulum.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

337867

Swiss Prot

Q8NBM4

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

PHGDHL1; UBAC2


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of various lysates using UBAC2 Rabbit pAb (A15977) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.

