

MUTYH Rabbit pAb

Catalog NO.: A1612

Basic Information

Observed MW

Calculated MW

60kDa

Category

Polyclonal Antibodies

Applications

IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Rat

Conjugation

Unconjugated

Background

This gene encodes a DNA glycosylase involved in oxidative DNA damage repair. The enzyme excises adenine bases from the DNA backbone at sites where adenine is inappropriately paired with guanine, cytosine, or 8-oxo-7,8-dihydroguanine, a major oxidatively damaged DNA lesion. The protein is localized to the nucleus and mitochondria. This gene product is thought to play a role in signaling apoptosis by the introduction of single-strand breaks following oxidative damage. Mutations in this gene result in heritable predisposition to colorectal cancer, termed MUTYH-associated polyposis (MAP). Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID	Swiss Prot
4595	Q9UIF7

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

MYH; MUTYH

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Immunofluorescence analysis of U2OS cells using MUTYH Rabbit pAb (A1612). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

