

# FADS1 Rabbit pAb

Catalog NO.: A16306 **1 Publications**

## Basic Information

### Observed MW

52kDa

### Calculated MW

52kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, both of which are characterized by conserved histidine motifs. This gene is clustered with family members FADS1 and FADS2 at 11q12-q13.1; this cluster is thought to have arisen evolutionarily from gene duplication based on its similar exon/intron organization.

## Recommended Dilutions

**WB** 1:500 - 1:2000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

| Gene ID | Swiss Prot |
|---------|------------|
| 3992    | O60427     |

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

D5D; TU12; FADS6; FADSD5; LLCDL1; FADS1


## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                       |                |                       |
|-------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                         | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                | IgG            | Affinity purification |
| <b>Storage</b>                                        |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.           |                |                       |
| Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. |                |                       |

Validation Data



Western blot analysis of various lysates using FADS1 Rabbit pAb (A16306) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 90s.

