FXN / Frataxin Rabbit pAb

Catalog NO.: A1745SP 2 Publications

Basic Information

Observed MW
14 kDa(mature FXN)/19 kDa(immature FXN)

Calculated MW
23 kDa/21 kDa/19 kDa

Category
Polyclonal Antibodies

Applications
WB, IF/ICC, IHC-P, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

This nuclear gene encodes a mitochondrial protein which belongs to the FRATAXIN family. The protein functions in regulating mitochondrial iron transport and respiration. The expansion of intronic trinucleotide repeat GAA from 8-33 repeats to >90 repeats results in Friedreich ataxia. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:2000 - 1:5000
IF/ICC	1:100 - 1:200
IHC-P	1:200 - 1:800

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (⩾1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
2395

Swiss Prot
Q16595

Immunogen
This information is considered to be commercially sensitive.

Synonyms
FA; X25; CyaY; FARR; FRDA; FXN / Frataxin

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.


Validation Data




Western blot analysis of various lysates using FXN / Frataxin Rabbit pAb (A1745SP) at 1:5000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 45 s.




Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using FXN / Frataxin Rabbit pAb (A1745SP) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using FXN / Frataxin Rabbit pAb (A1745SP) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of A549 cells using FXN / Frataxin Rabbit pAb (A1745SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunofluorescence analysis of Hep G2 cells using FXN / Frataxin Rabbit pAb (A1745SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.