SLC35D3 Rabbit pAb

Catalog NO.: A17880

Basic Information

Observed MW
44kDa

Calculated MW
44kDa

Category
Polyclonal Antibodies

Applications
WB, ELISA

Cross-Reactivity
Human

Conjugation
Unconjugated

Background

Predicted to enable antiporter activity and pyrimidine nucleotide-sugar transmembrane transporter activity. Predicted to be involved in carbohydrate transport and pyrimidine nucleotide-sugar transmembrane transport. Predicted to act upstream of or within energy homeostasis and positive regulation of protein exit from endoplasmic reticulum. Predicted to be located in early endosome and endoplasmic reticulum. Predicted to be integral component of membrane. Predicted to be active in Golgi apparatus.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
340146

Swiss Prot
Q5M8T2

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
FRCL1; SLC35D3

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using SLC35D3 Rabbit pAb (A17880) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 90s.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.