

# DTX3L Rabbit pAb

Catalog NO.: A18551 **1 Publications**

## Basic Information

### Observed MW

84kDa

### Calculated MW

84kDa

### Category

Polyclonal Antibodies

### Applications

WB, IF/ICC, ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

Enables several functions, including STAT family protein binding activity; ubiquitin-like protein ligase binding activity; and ubiquitin-protein transferase activity. Involved in several processes, including positive regulation of macromolecule metabolic process; positive regulation of protein localization; and protein ubiquitination. Located in cytosol; lysosome; and nucleoplasm. Part of protein-containing complex. Colocalizes with early endosome and nucleus.

## Recommended Dilutions

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

151636

### Swiss Prot

Q8TDB6

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

BBAP; RNF143; DTX3L

## Contact

## Product Information

|   |  |                      |
|---|--|----------------------|
| ☒ |  | order@abclonal.com   |
| ☒ |  | service@abclonal.com |
| ☒ |  | www.abclonal.com     |

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

**Validation Data**



Western blot analysis of lysates from Mouse thymus, using DTX3L Rabbit pAb (A18551) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.

