

MPC1 Rabbit pAb

Catalog NO.: A20195 **3 Publications**

Basic Information

Observed MW

12kDa

Calculated MW

12kDa

Category

Polyclonal Antibodies

Applications

WB, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

The protein encoded by this gene is part of an MPC1/MPC2 heterodimer that is responsible for transporting pyruvate into mitochondria. The encoded protein is found in the inner mitochondrial membrane. Defects in this gene are a cause of mitochondrial pyruvate carrier deficiency. Several transcript variants, some protein coding and one non-protein coding, have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:1000

IP

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

51660

Swiss Prot

Q9Y5U8

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MPYCD; BRP44L; CGI-129; SLC54A1; MPC1

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates using MPC1 Rabbit pAb (A20195) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.



Immunoprecipitation analysis of 300 µg extracts of K-562 cells using 3 µg MPC1 antibody (A20195). Western blot was performed from the immunoprecipitate using MPC1 antibody (A20195) at a dilution of 1:1000.

