TNKS1BP1 Rabbit pAb

Catalog NO.: A20388

Basic Information

Observed MW
Refer to figures

Calculated MW
182kDa

Category
Polyclonal Antibodies

Applications
WB, IHC-P, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

Enables ankyrin repeat binding activity and enzyme binding activity. Involved in cellular response to ionizing radiation; double-strand break repair; and positive regulation of protein phosphorylation. Located in several cellular components, including actin cytoskeleton; adherens junction; and heterochromatin. Part of CCR4-NOT complex.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
85456

Swiss Prot
Q9C0C2

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
TAB182; TNKS1BP1

Contact

Product Information

☒ | order@abclonal.com
☒ | service@abclonal.com
☒ | www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Human colon using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat stomach using TNKS1BP1 Rabbit pAb (A20388) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.