

SIGLEC10 Rabbit pAb

Catalog NO.: A20413

Basic Information

Observed MW

100kDa

Calculated MW

77kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Human, Rat

Conjugation

Unconjugated

Background

SIGLECs are members of the immunoglobulin superfamily that are expressed on the cell surface. Most SIGLECs have 1 or more cytoplasmic immune receptor tyrosine-based inhibitory motifs, or ITIMs. SIGLECs are typically expressed on cells of the innate immune system, with the exception of the B-cell expressed SIGLEC6 (MIM 604405).

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

89790

Swiss Prot

Q96LC7

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

SLG2; PRO940; SIGLEC-10; SIGLEC10

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from Raji cells, using SIGLEC10 Rabbit pAb (A20413) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Western blot analysis of lysates from Rat lung, using SIGLEC10 Rabbit pAb (A20413) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.

