BRG1/SMARCA4 Rabbit pAb

Catalog NO.: A2117SP **3 Publications**

Basic Information

Observed MW
220 kDa

Calculated MW
185 kDa/182 kDa/181 kDa

Category
Polyclonal Antibodies

Applications
WB, IF/ICC, IHC-P, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

The protein encoded by this gene is a member of the SWI/SNF family of proteins and is similar to the brahma protein of Drosophila. Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Mutations in this gene cause rhabdoid tumor predisposition syndrome type 2. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:1000 - 1:3000
IF/ICC	1:100 - 1:400
IHC-P	1:100 - 1:400

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (⩾1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
6597

Swiss Prot
P51532

Immunogen
This information is considered to be commercially sensitive.

Synonyms
BRG1; CSS4; SNF2; SWI2; MRD16; RTPS2; BAF190; SNF2L4; SNF2LB; hSNF2b; BAF190A; SNF2-beta; A4

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.


Validation Data



Western blot analysis of lysates from HeLa cells using BRG1/SMARCA4 Rabbit pAb (A2117SP) at 1:1000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1 s.



Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using BRG1/SMARCA4 Rabbit pAb (A2117SP) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of NIH/3T3 cells using BRG1/SMARCA4 Rabbit pAb (A2117SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS037, dilution 1:800) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.