

ABflo® 647 Rabbit anti-Human CD30 mAb

Catalog NO.: A22215

Basic Information

Observed MW

Calculated MW

14kDa/50kDa/63kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Human

Conjugation

ABflo® 647. Ex:648nm. Em:664nm.

Clone number

ARC52955

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

943

Swiss Prot

P28908

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD30; Ki-1; D1S166E


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Flow cytometry: 1X10^6 Human PBMC (untreated,left) and Human PBMC (treated with 5 µg/mL PHA for 72 hours,right) were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD3 mAb (A27177,5 µl/Test) and ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Human CD30 mAb (A22215,5 µl/Test,right). Cells in the lymphocyte gate were used for analysis.

