GNA13 Rabbit pAb

Catalog NO.: A24856

Basic Information

Observed MW
Refer to figures

Calculated MW
33kDa/44kDa

Category
Polyclonal Antibodies

Applications
IHC-P

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

Predicted to enable D5 dopamine receptor binding activity; G-protein beta/gamma-subunit complex binding activity; and GTPase activity. Predicted to be involved in several processes, including Rho protein signal transduction; activation of phospholipase D activity; and multicellular organism aging. Predicted to act upstream of or within several processes, including branching involved in blood vessel morphogenesis; negative regulation of vascular associated smooth muscle cell migration; and negative regulation of vascular associated smooth muscle cell proliferation. Located in cytosol and nucleus.

Recommended Dilutions

IHC-P	1:50 - 1:200

Immunogen Information

Gene ID
10672

Swiss Prot
Q14344

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
GNA13; G13; G protein subunit alpha 13

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Human liver tissue using GNA13 Rabbit pAb (A24856) at a dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using GNA13 Rabbit pAb (A24856) at a dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney tissue using GNA13 Rabbit pAb (A24856) at a dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.