

CXCL1 Rabbit mAb

Catalog NO.: A25014 **Recombinant**

Basic Information

Observed MW

45kDa (Recombinant Protein)

Calculated MW

10kDa

Category

Monoclonal Antibodies

Applications

WB, ELISA

Cross-Reactivity

Mouse

Conjugation

Unconjugated

Clone number

ARC65165

Background

This gene encodes a protein that is a member of the CXC subfamily of chemokines. Chemokines, which recruit and activate leukocytes, are classified by function (inflammatory or homeostatic) or by structure. This secretory protein is proposed to bind the G-protein coupled receptor chemokine (C-X-C motif) receptor 2 to recruit neutrophils. In mouse, deficiency of this gene is associated with colitis and with defects in immune cell recruitment to the lung.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID	Swiss Prot
14825	P12850

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

KC; Fsp; N51; gro; Gro1; Mgsa; Scyb1; CXCL1

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of recombinant mouse CXCL1/GRO-alpha Protein using CXCL1 Rabbit mAb (A25014) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 5ng/1ng per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.

