

ABflo® 647 Rabbit anti-Mouse CD14 mAb

Catalog NO.: A25191

Basic Information

Observed MW

Refer to figures

Calculated MW

39kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Mouse

Conjugation

ABflo® 647. Ex:648nm. Em:664nm.

Clone number

ARC66543

Background

This gene encodes a protein that plays an important role in the innate immune response and is expressed in monocyte/macrophage cells. This gene product acts as a co-receptor that binds several microbial and fungal molecules, including lipopolysaccharides (LPS). This proteins LPS-binding activity is enhanced by the LPS binding protein (LBP) to allow binding to the TLR4-MD-2 co-receptor complex. The product of this gene is found in two forms, either as a soluble protein or attached to the cell surface by a glycosylphosphatidylinositol anchor.

Recommended Dilutions

FC	5 µl per 10 ⁶ cells in 100 µl volume
----	---

Immunogen Information

Gene ID

12475

Swiss Prot

P10810

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD14


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		



Validation Data



Flow cytometry: 1X10^6 Neuro-2a cells (negative control,left) and RAW 264.7 cells (right) were surface-stained with ABflo® 647 Rabbit anti-Mouse CD14 mAb (A25191,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10^6 RAW 264.7 cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Mouse CD14 mAb (A25191,5 µl/Test,right).



Flow cytometry: 1X10⁶ C57BL/6 Mouse Bone Marrow-Derived M0 Macrophages were surface-stained with ABflo® 488 Rat anti-Mouse F4/80 mAb (A27479, 5 µl/Test) and ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, left) or ABflo® 647 Rabbit anti-Mouse CD14 mAb (A25191, 5 µl/Test, right).