

ABflo® 647 Rabbit anti-Mouse CD72 mAb

Catalog NO.: A25519

Basic Information

Observed MW

Refer to figures

Calculated MW

40kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Mouse

Conjugation

ABflo® 647. Ex:648nm. Em:664nm.

Clone number

ARC67530

Background

Predicted to enable carbohydrate binding activity and transmembrane signaling receptor activity. Predicted to be located in membrane. Predicted to be integral component of membrane. Predicted to be integral component of plasma membrane. Is expressed in brain; cerebellum external granule cell layer; cerebellum internal granule cell layer; hippocampus; and neocortex. Orthologous to human CD72 (CD72 molecule).

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

12517

Swiss Prot

P21855

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD72c; Ly-19; Ly-32; Lyb-2; Ly-m19


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		


Validation Data



Flow cytometry: 1X10⁶ NIH/3T3 cells (negative control,left) and C57BL/6 mouse splenocytes cells (right) were surface-stained with ABflo[®] 647 Rabbit anti-Mouse CD72 mAb (A25519,5 µl/Test,orange line) or ABflo[®] 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry: 1X10⁶ C57BL/6 mouse splenocytes cells were surface-stained with ABflo[®] 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo[®] 647 Rabbit anti-Mouse CD72 mAb (A25519,5 µl/Test,right).



Flow cytometry: 1×10^6 293T cells (negative control, left) and 293T (Transfection, right) cells were surface-stained with ABflo® 647 Rabbit anti-Mouse CD72 mAb (A25519, 5 μ l/Test, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Validation Data



Flow cytometry: 1×10^6 293T (Transfection) cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 μ l/Test, left) or ABflo® 647 Rabbit anti-Mouse CD72 mAb (A25519, 5 μ l/Test, right).

