

ABflo® 488 Rabbit anti-Mouse CD6 mAb

Catalog NO.: A25563

Basic Information

Observed MW

Calculated MW

64kDa/68kDa/72kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Mouse

Conjugation

ABflo® 488. Ex:491nm. Em:516nm.

Clone number

ARC66616

Background

Predicted to enable several functions, including identical protein binding activity; lipopolysaccharide binding activity; and lipoteichoic acid binding activity. Predicted to be involved in several processes, including heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules; lipopolysaccharide-mediated signaling pathway; and positive regulation of macromolecule metabolic process. Predicted to act upstream of or within cell adhesion. Predicted to be located in immunological synapse. Predicted to be spanning component of plasma membrane. Predicted to be part of T cell receptor complex. Predicted to be active in external side of plasma membrane. Orthologous to human CD6 (CD6 molecule).

Recommended Dilutions

FC 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

12511

Swiss Prot

Q61003

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Flow cytometry: 1X10⁶ NIH/3T3 cells (negative control,left) and C57BL/6 mouse Splenocytes (right) were surface-stained with ABflo[®] 488 Rabbit anti-Mouse CD6 mAb (A25563,5 µl/Test,orange line) or ABflo[®] 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10⁶ C57BL/6 mouse Splenocytes were surface-stained with ABflo[®] 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo[®] 488 Rabbit anti-Mouse CD6 mAb (A25563,5 µl/Test,right).

