D-Lactic acid-Histone H3-K9 Rabbit pAb

Catalog NO.: A26008

Basic Information

Observed MW

Calculated MW
15kDa

Category
Polyclonal Antibodies

Applications
DB, ELISA

Cross-Reactivity
Human

Conjugation
Unconjugated

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

DB	1:500 - 1:1000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8290/8350

Swiss Prot
Q16695/P68431

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.