

APC Rabbit anti-Human/Monkey Ki67 mAb

Catalog NO.: A27946

Basic Information

Observed MW

Calculated MW

359kDa

Category

Monoclonal Antibodies

Applications

FC (intra)

Cross-Reactivity

Human, Cynomolgus

Conjugation

APC. Ex:650nm. Em:660nm.

Clone number

ARC57564

Background

Enables protein C-terminus binding activity. Involved in regulation of chromosome segregation and regulation of mitotic nuclear division. Located in chromosome; nuclear body; and nucleolus. Colocalizes with condensed chromosome. Implicated in Crohn's disease; breast cancer; human immunodeficiency virus infectious disease; and pancreatic cancer. Biomarker of several diseases, including Barrett's esophagus; autoimmune disease of musculoskeletal system (multiple); endocrine gland cancer (multiple); gastrointestinal system cancer (multiple); and interstitial cystitis.

Recommended Dilutions

FC (intra) 5 µl per 10⁶ cells in 100 µl volume

Immunogen Information

Gene ID

Hu 4288 /Cyno 102135895

Swiss Prot

P46013

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

KIA; MIB-; MIB-1; PPP1R105


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.		

Validation Data



Flow cytometry: 1X10^6 knockout (KO) HeLa cells (negative control,left) and HeLa cells (right) were intracellularly-stained with APC Rabbit anti-Human/Monkey Ki67 mAb (A27946,5 µl/Test).



Flow cytometry: 1X10^6 Human PBMC (untreated,left) and Human PBMC (treated with 5 µg/mL PHA for 72 hours,right) were intracellularly-stained with ABflo® 450 Rabbit anti-Human cCD3 mAb (A27836,5 µl/Test) and APC Rabbit anti-Human/Monkey Ki67 mAb (A27946,5 µl/Test). Cells in the lymphocyte gate were used for analysis.

