

hCG beta Rabbit mAb

Catalog NO.: A27978 **Recombinant**

Basic Information

Observed MW

18kDa(non-glycosylated),20-35kDa(glycosylated)

Calculated MW

18kDa

Category

Monoclonal Antibodies

Applications

WB, IF-P, IHC-P, ELISA

Cross-Reactivity

Human

Conjugation

Unconjugated

Clone number

ARC75875

Background

This gene is a member of the glycoprotein hormone beta chain family and encodes the beta 5 subunit of chorionic gonadotropin (CG). Glycoprotein hormones are heterodimers consisting of a common alpha subunit and an unique beta subunit which confers biological specificity. CG is produced by the trophoblastic cells of the placenta and stimulates the ovaries to synthesize the steroids that are essential for the maintenance of pregnancy. The beta subunit of CG is encoded by 6 genes which are arranged in tandem and inverted pairs on chromosome 19q13.3 and contiguous with the luteinizing hormone beta subunit gene.

Recommended Dilutions

WB 1:2500 - 1:5000

IF-P 1:200 - 1:400

IHC-P 1:200 - 1:800

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

93659

Swiss Prot

P0DN86

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CGB; HCG; hCGB


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.09% Sodium azide,0.05% BSA,50% glycerol,pH7.3.		

Validation Data



Western blot analysis of lysates from BeWo cells using hCG beta Rabbit mAb (A27978) at 5000 dilution incubated overnight at 4°C. BeWo cells were treated with PNGase (6 U/μL) at 37°C for 30 minutes.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 μg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20 s.



Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using hCG beta Rabbit mAb (A27978) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Human placenta tissue using hCG beta Rabbit mAb (A27978, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

