

Serotonin transporter Rabbit pAb

Catalog NO.: A28369

Basic Information

Observed MW

83 kDa

Calculated MW

70 kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Mouse, Rat

Conjugation

Unconjugated

Background

Enables neurotransmitter transmembrane transporter activity; nitric-oxide synthase binding activity; and serotonin:sodium:chloride symporter activity. Involved in several processes, including nervous system development; platelet aggregation; and serotonin uptake. Located in focal adhesion and synapse. Is expressed in several structures, including early conceptus; heart; liver; nervous system; and ovary. Used to study autism spectrum disorder and sudden infant death syndrome. Human ortholog(s) of this gene implicated in several diseases, including alcohol dependence; anxiety disorder (multiple); depressive disorder (multiple); heart disease (multiple); and substance abuse (multiple). Orthologous to human SLC6A4 (solute carrier family 6 member 4).

Recommended Dilutions

WB 1:1000 - 1:2000

IHC-P 1:100 - 1:400

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000), a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID	Swiss Prot
15567	Q60857

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

Htt; Sert; 5-HTT

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of lysates from Rat brain using Serotonin transporter Rabbit pAb (A28369) at 1:1000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30 s.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Serotonin transporter Rabbit pAb (A28369) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Serotonin transporter Rabbit pAb (A28369) at a dilution of 1:300 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

