

APC anti-Mouse/Rat CD29/Integrin-β1 mAb

Catalog NO.: A28620

Basic Information

Observed MW

Calculated MW

88 kDa

Category

Monoclonal Antibodies

Applications

FC

Cross-Reactivity

Mouse,Rat

Conjugation

APC. Ex:650nm. Em:660nm.

Clone number

ACC0005

Background

CD29 is a 130 kDa protein, also known as integrin beta 1, the VLA-beta chain, or GPIIa. As a member of the integrin family, it is widely expressed on leukocytes, endothelial cells, smooth muscle, and epithelial cells. CD29 associates with CD49a-f to form the VLA-1 through VLA-6 complexes, respectively. It plays a crucial role in cell-cell and cell-matrix interactions. The HMβ1-1 antibody reacts with both mouse and rat CD29. It is capable of blocking cell adhesion and inhibiting T-cell proliferation.

Recommended Dilutions

FC ≤0.5 μg per million cells in 100 μl volume

Immunogen Information

Gene ID

16412,24511

Swiss Prot

P09055,P49134

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

CD29; Fnrb; gpIIa; Gm9863; 4633401G24Rik


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Armenian Hamster	IgG	Affinity purification
Storage		
Store at 2-8°C. Avoid freeze.		
Buffer: PBS with 0.09% Sodium azide,0.2% BSA,pH7.3.		

Validation Data



Flow cytometry:1X10^6 C57BL/6 mouse bone marrow cells were surface-stained with APC anti-Mouse/Rat CD29/Integrin-β1 mAb (A28620,0.5 μg,orange line) or APC Armenian Hamster IgG isotype control (5 μl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:1X10^6 SD rat bone marrow cells were surface-stained with APC anti-Mouse/Rat CD29/Integrin-β1 mAb (A28620,0.5 μg,orange line) or APC Armenian Hamster IgG isotype control (5 μl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

