

Neutrophil Elastase (ELANE) Rabbit mAb

Catalog NO.: A28985 **Recombinant**

Basic Information

Observed MW

29 kDa

Calculated MW

29 kDa

Category

Monoclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Clone number

ARC3858

Background

This gene encodes a member of the chymotrypsin-like family of serine protease enzymes that hydrolyzes a broad range of protein substrates including elastin. This gene is expressed by neutrophils where the encoded enzyme is stored in azurophil granules. Upon neutrophil activation, the active enzyme is released into the extracellular mileu. Mice lacking the encoded protein exhibit increased susceptibility to sepsis and death following intraperitoneal infection with Gram negative bacteria. This gene is located adjacent to a related proteinase gene on chromosome 10.

Recommended Dilutions

WB 1:2000 - 1:10000

IHC-P 1:100 - 1:400

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000), a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID	Swiss Prot
50701	Q3UP87

Immunogen

This information is considered to be commercially sensitive.

Synonyms

NE; Ela2; F430011M15Rik


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS with 0.09% sodium azide,0.05% BSA,50% glycerol,pH7.3		

Validation Data



Western blot analysis of various lysates using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at 1:5000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Negative control (NC): Rat kidney.

Exposure time: 90 s.



Western blot analysis of lysates from Mouse bone marrow using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at 1:5000 dilution incubated overnight at 4°C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 1 s.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a dilution of



Immunohistochemistry analysis of paraffin-embedded Human esophagus tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a dilution of




Immunohistochemistry analysis of paraffin-embedded Mouse bone marrow tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a

1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.


1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.


Validation Data



Immunohistochemistry analysis of paraffin-embedded Mouse spleen tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat bone marrow tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue using Neutrophil Elastase (ELANE) Rabbit mAb (A28985) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

