

# FKBP51/FKBP5 Rabbit pAb

Catalog NO.: A3863 **2 Publications**

## Basic Information

### Observed MW

51kDa

### Calculated MW

51kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds to the immunosuppressants FK506 and rapamycin. It is thought to mediate calcineurin inhibition. It also interacts functionally with mature hetero-oligomeric progesterone receptor complexes along with the 90 kDa heat shock protein and P23 protein. This gene has been found to have multiple polyadenylation sites. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

**WB** 1:200 - 1:2000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

| Gene ID | Swiss Prot |
|---------|------------|
| 2289    | Q13451     |

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

P54; AIG6; FKBP51; FKBP54; PPIase; Ptg-10; FKBP51/FKBP5

## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of various lysates using FKBP51/FKBP5 Rabbit pAb (A3863) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.

