

Histone H3.3 Rabbit mAb

Catalog NO.: A4835 **Recombinant** **2 Publications**

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

Monoclonal Antibodies

Applications

WB, IHC-P, ELISA, ChIP, ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Conjugation

Unconjugated

Clone number

ARC0255

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene contains introns and its mRNA is polyadenylated, unlike most histone genes. The protein encoded is a replication-independent member of the histone H3 family.

Recommended Dilutions

WB 1:1000 - 1:6000

IHC-P 1:100 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

ChIP 2µg antibody for 5µg-10µg of Chromatin

ChIP-seq 1:50 - 1:200

Immunogen Information

Gene ID	Swiss Prot
3020	P84243

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3F3; H3-3B; H3.3A; H3F3A; BRYLIB1; Histone H3.3

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from Rat lung, using Histone H3.3 Rabbit mAb (A4835) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates using Histone H3.3 Rabbit mAb (A4835) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Histone H3.3 antibody (A4835) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Rat kidney using Histone H3.3 Rabbit mAb (A4835) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human esophageal using Histone H3.3 Rabbit mAb (A4835) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse liver using Histone H3.3 Rabbit mAb (A4835) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and Histone H3.3 Rabbit mAb (A4835). The ChIP sequencing results indicate the enrichment pattern of Histone H3.3 in selected genomic region and representative gene loci (GAPDH), as shown in figure.

