

SRRM1 Rabbit pAb

Catalog NO.: A6066 **1 Publications**

Basic Information

Observed MW

102kDa

Calculated MW

102kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse

Conjugation

Unconjugated

Background

Enables RNA binding activity. Predicted to be involved in mRNA splicing, via spliceosome. Located in nuclear speck. Part of catalytic step 2 spliceosome. Biomarker of gestational diabetes.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:100 - 1:500

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

10250

Swiss Prot

Q8IYB3

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

160-KD; POP101; SRM160; SRRM1

Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.


Validation Data



Western blot analysis of various lysates using SRRM1 Rabbit pAb (A6066) at 1:3000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human esophageal cancer using SRRM1 Rabbit pAb (A6066) at dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using SRRM1 Rabbit pAb (A6066) at dilution of 1:400 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.

