

DARS Rabbit pAb

Catalog NO.: A6574 **1 Publications**

Basic Information

Observed MW

57kDa

Calculated MW

57kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, IP, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

This gene encodes a member of a multienzyme complex that functions in mediating the attachment of amino acids to their cognate tRNAs. The encoded protein ligates L-aspartate to tRNA(Asp). Mutations in this gene have been found in patients showing hypomyelination with brainstem and spinal cord involvement and leg spasticity. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:50 - 1:200

IP
0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

ELISA
Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Contact

Immunogen Information

Gene ID	Swiss Prot
1615	P14868

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.


Synonyms
DARS; HBSL; aspRS

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source Rabbit	Isotype IgG	Purification Affinity purification
Storage Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Western blot analysis of various lysates using DARS Rabbit pAb (A6574) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.




Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunoprecipitation analysis of 200 µg extracts of 293T cells using 1 µg DARS antibody (A6574). Western blot was performed from the immunoprecipitate using DARS antibody (A6574) at a dilution of 1:1000.



Immunohistochemistry analysis of paraffin-embedded Rat brain using DARS Rabbit pAb (A6574) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Mouse spinal cord using DARS Rabbit pAb (A6574) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Mouse brain using DARS Rabbit pAb (A6574) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.