

# BCL6 Rabbit pAb

Catalog NO.: A7173 **4 Publications**

## Basic Information

### Observed MW

86kDa

### Calculated MW

79kDa

### Category

Polyclonal Antibodies

### Applications

WB, IF/ICC, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### Conjugation

Unconjugated

## Background

The protein encoded by this gene is a zinc finger transcription factor and contains an N-terminal POZ domain. This protein acts as a sequence-specific repressor of transcription, and has been shown to modulate the transcription of STAT-dependent IL-4 responses of B cells. This protein can interact with a variety of POZ-containing proteins that function as transcription corepressors. This gene is found to be frequently translocated and hypermutated in diffuse large-cell lymphoma (DLCL), and may be involved in the pathogenesis of DLCL. Alternatively spliced transcript variants encoding different protein isoforms have been found for this gene.

## Recommended Dilutions

**WB** 1:500 - 1:1000

**IF/ICC** 1:50 - 1:100

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

| Gene ID | Swiss Prot |
|---------|------------|
| 604     | P41182     |

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

BCL5; LAZ3; BCL6A; ZNF51; ZBTB27; BCL6

## Contact

## Product Information

|   |  |                      |
|---|--|----------------------|
| ☒ |  | order@abclonal.com   |
| ☒ |  | service@abclonal.com |
| ☒ |  | www.abclonal.com     |

## Source

Rabbit

## Isotype

IgG

## Purification


Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from Raji cells, using BCL6 Rabbit pAb (A7173) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.



Immunofluorescence analysis of A549 cells using BCL6

Rabbit pAb (A7173).Secondary antibody: Cy3-conjugated

Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution.

