YTHDC1 Rabbit pAb

Catalog NO.: A7318SP **3 Publications**

Basic Information

Observed MW
110 kDa

Calculated MW
85 kDa/83 kDa

Category
Polyclonal Antibodies

Applications
WB, IP, IHC-P, ELISA

Cross-Reactivity
Human, Mouse, Rat

Conjugation
Unconjugated

Background

Enables N6-methyladenosine-containing RNA binding activity. Involved in mRNA export from nucleus; mRNA splice site selection; and regulation of gene expression. Located in nuclear speck and plasma membrane.

Recommended Dilutions

WB 1:1000 - 1:3000

IP
0.5 μg - 4 μg antibody for 200 μg - 400 μg extracts of whole cells

IHC-P 1:1000 - 1:4000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (⩾1:10000)

Immunogen Information

Gene ID
91746

Swiss Prot
Q96MU7

Immunogen
This information is considered to be commercially sensitive.

Synonyms
YT521; YT521-B; YTHDC1

Contact

Product Information

order@abclonal.com
service@abclonal.com
www.abclonal.com


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.


Validation Data




Western blot analysis of lysates from MOLT-4 cells using YTHDC1 Rabbit pAb (A7318SP) at 1:1000 dilution incubated overnight at 4°C.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 0.5 s.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.





Immunohistochemistry analysis of paraffin-embedded Human pancreas tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.








Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Rat skin tissue using YTHDC1 Rabbit pAb (A7318SP) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.