

NMDAR1 Rabbit pAb

Catalog NO.: A7677SP **16 Publications**

Basic Information

Observed MW

120 kDa

Calculated MW

99-108 kDa

Category

Polyclonal Antibodies

Applications

WB, IF-P, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Background

The protein encoded by this gene is a critical subunit of N-methyl-D-aspartate receptors, members of the glutamate receptor channel superfamily which are heteromeric protein complexes with multiple subunits arranged to form a ligand-gated ion channel. These subunits play a key role in the plasticity of synapses, which is believed to underlie memory and learning. Cell-specific factors are thought to control expression of different isoforms, possibly contributing to the functional diversity of the subunits. Alternatively spliced transcript variants have been described.

Recommended Dilutions

WB 1:5000 - 1:20000

IF-P 1:200 - 1:400

IHC-P 1:200 - 1:800

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements. For high-ratio antibody dilutions (≥1:10000)

Immunogen Information

Gene ID	Swiss Prot
2902	Q05586

Immunogen

This information is considered to be commercially sensitive.

Synonyms

NR1; MRD8; GluN1; NMDA1; DEE101; NDHMSD; NDHMSR; NMD-R1; NMDAR1


Contact

Product Information

☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS, pH 7.3, containing 50% glycerol. Preserved with Proclin300 or sodium azide. May contain 0.05% BSA as specified on the Certificate of Analysis.		

Validation Data



Western blot analysis of various lysates using NMDAR1 Rabbit pAb (A7677SP) at 1:10000 dilution incubated overnight at 4°C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Negative control (NC): Mouse liver, Rat liver.


Exposure time: 30 s.





Immunohistochemistry analysis of paraffin-embedded Human brain tissue using NMDAR1 Rabbit pAb (A7677SP) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using NMDAR1 Rabbit pAb (A7677SP) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using NMDAR1 Rabbit pAb (A7677SP) at a dilution of 1:500 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Mouse brain tissue using NMDAR1 Rabbit pAb (A7677SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

Antibody | Protein | ELISA Kit | Enzyme | NGS | Service

Confocal imaging of paraffin-embedded Rat brain tissue using NMDAR1 Rabbit pAb (A7677SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit <https://abclonal.com> for a complete list of recommended products.