

# POMP Rabbit pAb

Catalog NO.: A9067 **1 Publications**

## Basic Information

### Observed MW

16kDa

### Calculated MW

16kDa

### Category

Polyclonal Antibodies

### Applications

WB, ELISA

### Cross-Reactivity

Human

### Conjugation

Unconjugated

## Background

The protein encoded by this gene is a molecular chaperone that binds 20S preproteasome components and is essential for 20S proteasome formation. The 20S proteasome is the proteolytically active component of the 26S proteasome complex. The encoded protein is degraded before the maturation of the 20S proteasome is complete. A variant in the 5' UTR of this gene has been associated with KLINK syndrome, a rare skin disorder.

## Recommended Dilutions

**WB** 1:500 - 1:1000

### ELISA

Recommended starting concentration is 1 µg/mL.  
Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

| Gene ID | Swiss Prot |
|---------|------------|
| 51371   | Q9Y244     |

### Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

### Synonyms

UMP1; PRAAS2; HSPC014; C13orf12; PNAS-110; POMP


## Contact

## Product Information

|                                     |  |                      |
|-------------------------------------|--|----------------------|
| <input checked="" type="checkbox"/> |  | order@abclonal.com   |
| <input checked="" type="checkbox"/> |  | service@abclonal.com |
| <input checked="" type="checkbox"/> |  | www.abclonal.com     |

|                                                       |                |                       |
|-------------------------------------------------------|----------------|-----------------------|
| <b>Source</b>                                         | <b>Isotype</b> | <b>Purification</b>   |
| Rabbit                                                | IgG            | Affinity purification |
| <b>Storage</b>                                        |                |                       |
| Store at -20°C. Avoid freeze / thaw cycles.           |                |                       |
| Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. |                |                       |

Validation Data



Western blot analysis of various lysates using POMP Rabbit pAb (A9067) at 1:3000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 90s.

