

eIF3B Rabbit mAb

Catalog NO.: A9143 **Recombinant** **1 Publications**

Basic Information

Observed MW

116kDa

Calculated MW

93kDa

Category

Monoclonal Antibodies

Applications

WB, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Clone number

ARC1448

Background

Enables RNA binding activity. Contributes to translation initiation factor activity. Involved in several processes, including IRES-dependent viral translational initiation; translational initiation; and viral translational termination-reinitiation. Located in extracellular exosome. Part of eukaryotic translation initiation factor 3 complex.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL.
Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID	Swiss Prot
8662	P55884

Immunogen

Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms

PRT1; EIF3S9; EIF3-ETA; EIF3-P110; EIF3-P116; eIF3B


Contact

Product Information


☒		order@abclonal.com
☒		service@abclonal.com
☒		www.abclonal.com

Source	Isotype	Purification
Rabbit	IgG	Affinity purification
Storage		
Store at -20°C. Avoid freeze / thaw cycles.		
Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.		

Validation Data



Western blot analysis of various lysates using eIF3B Rabbit mAb (A9143) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 1s.



Confocal imaging of NIH/3T3 cells using eIF3B Rabbit mAb (A9143, dilution 1:100) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

Confocal imaging of C6 cells using eIF3B Rabbit mAb (A9143, dilution 1:100) followed by a further incubation with Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

