Phospho-Beclin 1-Ser93 Rabbit pAb

Catalog NO.: AP1287 **1 Publications**

Basic Information

Observed MW
60kDa

Calculated MW
52kDa

Category
Phosphorylated Antibodies

Applications
WB, ELISA

Cross-Reactivity
Human

Conjugation
Unconjugated

Background

This gene encodes a protein that regulates autophagy, a catabolic process of degradation induced by starvation. The encoded protein is a component of the phosphatidylinositol-3-kinase (PI3K) complex which mediates vesicle-trafficking processes. This protein is thought to play a role in multiple cellular processes, including tumorigenesis, neurodegeneration and apoptosis. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:100 - 1:500

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8678

Swiss Prot
Q14457

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
ATG6; VPS30; beclin1; Phospho-Beclin 1-Ser93

Contact

Product Information

☒ | order@abclonal.com

☒ | service@abclonal.com

☒ | www.abclonal.com


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20°C. Avoid freeze / thaw cycles.
Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using Phospho-Beclin 1-Ser93 Rabbit pAb (AP1287) at 1:500 dilution. HeLa cells were treated with Calyculin A (100 nM) at 37°C for 30 minutes.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 180s.

For research use only. Not for therapeutic or diagnostic purposes.

Please visit **https://abclonal.com** for a complete list of recommended products.