

H2-Aa Rabbit pAb

Catalog No.: A18325**2 Publications**

Basic Information

Observed MW

28 kDa

Calculated MW

28 kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross Reactivity

Human, Mouse, Rat

Background

Enables peptide antigen binding activity. Involved in response to interferon-gamma. Acts upstream of or within antigen processing and presentation of exogenous peptide antigen via MHC class II and positive regulation of T cell differentiation. Located in external side of plasma membrane and lysosome. Part of MHC class II protein complex. Is expressed in central nervous system; retina; and thymus primordium. Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); gastrointestinal system cancer (multiple); glomerulonephritis (multiple); hematologic cancer (multiple); and systemic scleroderma (multiple). Orthologous to several human genes including HLA-DQA1 (major histocompatibility complex, class II, DQ alpha 1).

Recommended Dilutions

WB 1:500 - 1:1000**IHC-P** 1:200 - 1:2000**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

14960

Swiss-Prot

P14434

Immunogen

This information is considered to be commercially sensitive.

Synonyms

Ia1; H2Aa; Ia-1; H-2Aa; Aalpha; IAalpha; I-Aalpha; H2-Aa

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purity

Affinity purification

Storage

Store at -20°C

Validation Data



Western blot analysis of various lysates using H2-Aa Rabbit pAb (A18325) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.