

TERF1 Rabbit pAb

Catalog No.: A0137**3 Publications**

Basic Information

Observed MW

50 kDa

Calculated MW

50 kDa

Category

Polyclonal Antibodies

Applications

WB, IHC-P, ELISA

Cross Reactivity

Human, Mouse, Rat

Background

This gene encodes a telomere specific protein which is a component of the telomere nucleoprotein complex. This protein is present at telomeres throughout the cell cycle and functions as an inhibitor of telomerase, acting in cis to limit the elongation of individual chromosome ends. The protein structure contains a C-terminal Myb motif, a dimerization domain near its N-terminus and an acidic N-terminus. Multiple transcripts of this gene are alternatively spliced products.

Recommended Dilutions

WB 1:500 - 1:2000**IHC-P** 1:50 - 1:200**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

7013

Swiss-Prot

P54274

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TRF; PIN2; TRF1; TRBF1; t-TRF1; hTRF1-AS; TERF1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG


Purity

Affinity purification


Storage

Store at -20°C

Validation Data



Immunohistochemistry analysis of paraffin-embedded Human gastric cancer using TERF1 Rabbit pAb (A0137) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Western blot analysis of various lysates, using TERF1 Rabbit pAb (A0137) at 1:2000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.