

CES2 Rabbit pAb**Catalog No.: A1514****Basic Information****Observed MW**

67 kDa

Calculated MW

62 kDa

Category

Polyclonal Antibodies

Applications

WB, IF/ICC, ELISA

Cross Reactivity

Human

Background

This gene encodes a member of the carboxylesterase large family. The family members are responsible for the hydrolysis or transesterification of various xenobiotics, such as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-brain barrier system. The protein encoded by this gene is the major intestinal enzyme and functions in intestine drug clearance. Alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions**WB** 1:500 – 1:2000**IF/ICC** 1:50 – 1:200**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information**Gene ID**

8824

Swiss-Prot

O00748

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

iCE; CE-2; PCE-2; CES2A1; CES2

Contact | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn**Product Information****Source**

Rabbit

Isotype

IgG


Purity

Affinity purification


Storage

Store at -20°C

Validation Data



Western blot analysis of lysates from SW620 cells using CES2 Rabbit pAb (A1514) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of MCF-7 cells using CES2 Rabbit pAb (A1514). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.