

FIG4 Rabbit pAb**Catalog No.: A15372****Basic Information****Observed MW**

104 kDa

Calculated MW

104 kDa

Category

Polyclonal Antibodies

Applications

WB, ELISA

Cross Reactivity

Human, Mouse

Background

The protein encoded by this gene belongs to the SAC domain-containing protein gene family. The SAC domain, approximately 400 amino acids in length and consisting of seven conserved motifs, has been shown to possess phosphoinositide phosphatase activity. The yeast homolog, Sac1p, is involved in the regulation of various phosphoinositides, and affects diverse cellular functions such as actin cytoskeleton organization, Golgi function, and maintenance of vacuole morphology. Membrane-bound phosphoinositides function as signaling molecules and play a key role in vesicle trafficking in eukaryotic cells. Mutations in this gene have been associated with Charcot-Marie-Tooth disease, type 4J.

Recommended Dilutions**WB** 1:500 - 1:2000**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information**Gene ID**

9896

Swiss-Prot

Q92562

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

BOP; YVS; BTOP; SAC3; ALS11; CMT4J; KIAA0274; dJ249I4.1; FIG4

Contact | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn**Product Information****Source**

Rabbit

Isotype

IgG


Purity

Affinity purification

Storage

Store at -20°C

Validation Data



Western blot analysis of various lysates using FIG4 Rabbit pAb (A15372) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.