Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb

Catalog No.: A17899

www.abclonal.com

Basic Information

Observed MW
17 kDa

Calculated MW
15 kDa

Category
Polyclonal Antibodies

Applications
WB, IF/ICC, IHC-P, DB, ELISA

Cross Reactivity
Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200
IHC-P	1:50 - 1:200
DB	1:500 - 1:1000

ELISA
Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8290/8350

Swiss-Prot
Q16695/P68431

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H3/A; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FA; H3C10; H3C11; H3C12; HIST1H3A; Asymmetric DiMethyl-Histone H3-R17

Contact

400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

Product Information


Source
Rabbit

Isotype
IgG


Purity
Affinity purification

Storage
Store at -20℃


Validation Data




Immunofluorescence analysis of NIH/3T3 cells using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A17899) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A17899) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Western blot analysis of various lysates using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A17899) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1: 10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 180s.



Dot-blot analysis of all sorts of peptides using Asymmetric DiMethyl-Histone H3-R17 Rabbit pAb (A17899) at 1:1000 dilution.