

DJ-1/PARK7 Rabbit mAb**Catalog No.: A19097**

Recombinant

3 Publications

Basic Information**Observed MW**

23 kDa

Calculated MW

20 kDa

Category

Monoclonal Antibodies

Applications

WB, IP, IF/ICC, IHC-P, ELISA

Cross Reactivity

Human, Mouse, Rat

Clone No.

ARC0354

Background

The product of this gene belongs to the peptidase C56 family of proteins. It acts as a positive regulator of androgen receptor-dependent transcription. It may also function as a redox-sensitive chaperone, as a sensor for oxidative stress, and it apparently protects neurons against oxidative stress and cell death. Defects in this gene are the cause of autosomal recessive early-onset Parkinson disease 7. Two transcript variants encoding the same protein have been identified for this gene.

Recommended Dilutions**WB** 1:1000 - 1:6000**IP**

0.5 µg - 4 µg antibody for 200 µg - 400 µg extracts of whole cells

IF/ICC 1:100 - 1:200**IHC-P** 1:200 - 1:800**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information**Gene ID**

11315

Swiss-Prot

Q99497

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

DJ1; DJ-1; GATD2; HEL-S-67p; DJ-1/PARK7

Contact | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn**Product Information****Source**

Rabbit

Isotype

IgG


Purity

Affinity purification


Storage

Store at -20°C


Validation Data




Immunofluorescence analysis of HeLa cells and DJ-1/PARK7 knockout (KO) HeLa cells using DJ-1/PARK7 Rabbit mAb (A19097) at a dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using DJ-1/PARK7 Rabbit mAb (A19097) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis tissue using DJ-1/PARK7 Rabbit mAb (A19097) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IHC staining.



Immunoprecipitation of DJ-1/PARK7 from 300 µg extracts of HeLa cells was performed using 2 µg of DJ-1/PARK7 Rabbit mAb (A19097). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1x Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using DJ-1/PARK7 Rabbit mAb (A19097) at a dilution of 1:1000.



Western blot analysis of various lysates using DJ-1/PARK7 Rabbit mAb (A19097) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.