

# Phospho-ACLY-S455 Rabbit pAb

**Catalog No.: AP0779****2 Publications**

## Basic Information

**Observed MW**

125 kDa

**Calculated MW**

121kDa

**Category**

Polyclonal Antibodies

**Applications**

WB, IP, IHC-P, ELISA

**Cross Reactivity**

Human, Mouse, Rat

## Background

ATP citrate lyase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. The enzyme is a tetramer (relative molecular weight approximately 440,000) of apparently identical subunits. It catalyzes the formation of acetyl-CoA and oxaloacetate from citrate and CoA with a concomitant hydrolysis of ATP to ADP and phosphate. The product, acetyl-CoA, serves several important biosynthetic pathways, including lipogenesis and cholesterologenesis. In nervous tissue, ATP citrate-lyase may be involved in the biosynthesis of acetylcholine. Multiple transcript variants encoding distinct isoforms have been identified for this gene.

## Recommended Dilutions

**WB** 1:500 - 1:2000**IP**

0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

**IHC-P** 1:50 - 1:100**ELISA**

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**

47

**Swiss-Prot**

P53396

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

ACL; ATPCL; CLATP; Phospho-ACLY-S455

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG


**Purity**

Affinity purification


**Storage**

Store at -20°C


## Validation Data




Immunoprecipitation analysis of 200 µg extracts of NIH/3T3 cells, using 3 µg Phospho-ACLY-S455 pAb (AP0779). Western blot was performed from the immunoprecipitate using Phospho-ACLY-S455 pAb (AP0779) at a dilution of 1:1000. NIH/3T3 cells were treated with Insulin (100 nM) at 37°C for 10 minutes after serum-starvation overnight.




Immunohistochemistry analysis of paraffin-embedded Rat pancreas using Phospho-ACLY-S455 Rabbit pAb (AP0779) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Phospho-ACLY-S455 Rabbit pAb (AP0779) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse pancreas using Phospho-ACLY-S455 Rabbit pAb (AP0779) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Western blot analysis of lysates from HeLa cells using Phospho-ACLY-S455 Rabbit pAb (AP0779) at 1:1000 dilution incubated overnight at 4°C. HeLa cells were treated with insulin (50 nM) at 37°C for 30 minutes.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60 s.



Western blot analysis of lysates from NIH/3T3 cells using Phospho-ACLY-S455 Rabbit pAb (AP0779) at 1:1000 dilution incubated overnight at 4°C. NIH/3T3 cells were treated with insulin (200 nM) at 37°C for 30 minutes.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60 s.



Western blot analysis of lysates from C6 cells using Phospho-ACLY-S455 Rabbit pAb (AP0779) at 1:1000 dilution incubated overnight at 4°C. C6 cells were treated with insulin (100 nM) at 37°C for 30 minutes.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60 s.