Recombinant Human B7-H1/PD-L1/CD274 Protein

Catalog No.: RP00068

www.abclonal.com

Sequence Information

Species
Human

Gene ID
29126

Swiss Prot
Q9NZQ7

Tags
C-His

Synonyms
B7-H, B7H1, PDL1, PD-L1, hPD-L1, PDCD1L1, PDCD1LG1,CD274, PDL1,B7H1,PD-L1,PDCD1L1, PDCD1LG1, B7-H, CD274 molecule

Product Information

Source
HEK293 cells

Purification
≥ 95 % as determined by SDS-PAGE;≥95 % as determined by HPLC.

Calculated MW
26.14 kDa

Observed MW
35-40 kDa

Endotoxin
< 0.1 EU/μg of the protein by LAL method.

Formulation
Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

This protein is an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation.

Basic Information

Description
Recombinant Human B7-H1/PD-L1/CD274 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Phe19-Thr239) of human PD-L1/B7-H1 (Accession #NP_054862.1) fused with a 6×His tag at the C-terminus.

Activity
1.Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human PD-L1 at 10 μg/mL (100 μL/well) can bind Recombinant Human PD-1 with a linear range of 0.25-1.02 μg/mL.
2.Measured by its binding ability in a functional ELISA.Immobilized PE anti-human CD274 (B7-H1, PD-L1) Antibody at 1μg/mL (25 μL/well) can bind Human CD274 with a linear range of 0.46-18ng/mL.

Shipping
Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage
Store at -20℃


Operational Notes
For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.


Validation Data




Immobilized Human PD-L1 (Catalog: RP00068) at 200ng/mL (100 µL/well) can bind PD-L1/CD274 Rabbit mAb (Catalog: A20270) with a linear range of 0.06-1.25 ng/mL.



The purity of Human PD-L1/B7-H1 Protein (Cat.RP00068) was greater than 95% as determined by SEC-HPLC.



Recombinant Human B7-H1/PD-L1/CD274 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Human PD-L1 His Tag (Catalog: RP00068) at 5ug/mL (100uL/well) can bind Recombinant Human PD-1 with a linear range of 0.16-1.09 µg /mL.