Recombinant Human LIMP II/SCARB2/CD36L2 Protein

Catalog No.: RP00086

Sequence Information

Species	Gene ID
Human	950
Swiss Prot	**Tags**
Q14108	C-hFc&His

Synonyms

AMRF, CD36L2, EPM4, HLGP85, LGP85, LIMP-2, LIMPII, SR-BII, SCARB2,CD36L2,EPM4,HLGP85, LGP85,LIMP-2,LIMPII,SR-BII

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE.
Calculated MW	**Observed MW**
73.33 kDa	90-110 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The protein encoded by this gene is a type III glycoprotein that is located primarily in limiting membranes of lysosomes and endosomes. Earlier studies in mice and rat suggested that this protein may participate in membrane transportation and the reorganization of endosomal/lysosomal compartment. The protein deficiency in mice was reported to impair cell membrane transport processes and cause pelvic junction obstruction, deafness, and peripheral neuropathy. Further studies in human showed that this protein is a ubiquitously expressed protein and that it is involved in the pathogenesis of HFMD (hand, foot, and mouth disease) caused by enterovirus-71 and possibly by coxsackievirus A16. Mutations in this gene caused an autosomal recessive progressive myoclonic epilepsy-4 (EPM4), also known as action myoclonus-renal failure syndrome (AMRF). Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Basic Information

Description

Recombinant Human LIMP II/SCARB2/CD36L2 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Arg27-Thr432) of human LIMPII/SR-B2 (Accession #NP_005497.1) fused with an Fc, 6×His tag at the C-terminus.

Activity

Measured by its binding ability in a functional ELISA. Immobilized Human LDLR/LDL Receptor at 4 μg/mL (100 μL/well) can bind Human SCARB2 with a linear range of 0.016-3.79 μg/mL.

Shipping

Lyophilized proteins are shipped at ambient temperature.
Liquid proteins are shipped at frozen temperature with dry ice.
Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20℃

Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.

Contact


400-999-6126
cn.market@abclonal.com.cn
www.abclonal.com.cn

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.

Validation Data



Recombinant Human LIMP II/SCARB2/CD36L2 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human LDLR/LDL Receptor at 4 μg/mL (100 μL/well) can bind Human SCARB2 with a linear range of 0.016-3.79 μg/mL.